

Steroids

1. Clobetasol propionate 0.05 % in orabase (5 gm/pack)
Sig. apply tid pc.



2. Dexamethasone mouthwash 0.05 % (300 ml/bottle)
Sig. use as mouthwash bid pc.



3. Fluocinolone acetonide 0.1 % in orabase (5 gm/pack)
Sig. apply tid pc.



**4. Fluocinonide 0.05 % in orabase (5 gm/pack)
Sig. apply tid pc.**



**5. Fluocinolone acetonide solution 0.1 % (10 ml/bottle)
Sig. apply tid pc.**



**6. Fluocinolone acetonide 0.1 % with clotrimazole 0.1 %
gel (5 gm/pack)
Sig. apply tid pc.**



**7. Triamcinolone acetonide 0.1 % in orabase (5 gm/pack)
Sig. apply tid pc.**

ใน 1 กรัมของเคนาลอกมี 1 มก. (0.1%) ไตรแอมซิโนโลน ในจีฟี่ง
สำหรับใช้ภายในปาก โดยเฉพาะที่ สำหรับรักษาแผลในช่องปาก จีฟี่ง
สำหรับทาในปาก ประกอบด้วยเฮลลาดิน เพกตินและโซเดียมคาร์บอ-
เนต เมธิลเซลลูโลส ทำให้ยึดติดอยู่ในปากได้ง่าย จึงได้ผลในการรักษา
และช่วยเคลือบแผล หรือบริเวณที่อักเสบด้วย

ข้อบ่งใช้ รักษาและบรรเทาอาการอักเสบในช่องปาก

ข้อห้ามใช้ ผู้ที่มีประวัติแพ้ยาหรือสารประกอบในยา
การอักเสบในปากหรือลำคอที่เกิดจากเชื้อรา, ไวรัส หรือแบคทีเรีย
คำเตือน การใช้ในหญิงมีครรภ์ ยังไม่มีการศึกษาถึงผลของยาที่มีต่อ
ทารกในครรภ์ ดังนั้นจึงไม่ควรใช้ในหญิงที่ตั้งครรภ์หรือคาดว่าจะมี
นอกจากได้รับคำแนะนำจากแพทย์หรือทันตแพทย์

ข้อควรระวัง ไม่ควรใช้ยาคอร์ติโคสเตอรอยด์ใดๆ ในผู้ที่เป็วันวัณโรค
โรคกระเพาะอาหารอักเสบ หรือโรคเบาหวาน, ปฏิกริยาของเชื้อปาก
อาจถูกกดด้วยยาแบบเฉพาะที่ ดังนั้นอาจทำให้ความรุนแรงของเชื้อ
โรคในปากเพิ่มขึ้น โดยที่ร่างกายไม่มีอาการ



**8. Triamcinolone acetonide mouthwash 0.1 % (120 ml/bottle)
Sig. use as mouthwash tid pc.**



MIMS 99th Edition 2005 page 371

**9. Prednisolone cream (Prednisil®) 5 mg/g (5 gm/pack)
(Prednisolone BP 0.5% in cream base)
Sig. apply od-bid pc.**

I: Atropic dermatitis: infantile & allergic eczema, neurodermatitis, pruritus with lichenification. Contact dermatitis & non-specific pruritus.

D: Apply once-bid.

CI: Viral diseases, bacterial & fungal skin infections, acne rosacea & perioral dermatitis.

SP: Prolonged use in childn.

AR: Local atropic changes(prolonged & intensive treatment).Skin collagen loss.

Hypercorticism. Itching, folliculitis, hypertrichosis, acneiform eruptions, hypopigmentation, perioral & allergic contact dermatitis, skin maceration, secondary infections, striae & miliaria.



10. Prednisil-N® cream


Sig. apply od-bid pc.

1 กรัม ประกอบด้วย เพรดนิโซโลน 5 มก. นีโอไมซิน ซัลเฟต 5 มก. เทียบเท่ากับ นีโอไมซิน 3.25 มก.

ข้อบ่งใช้ สำหรับใช้ในโรคผิวหนังผื่นคันตามแก้ม และร่องจมูก ในเด็กเล็กๆ (infantile eczema) โรคผิวหนังผื่นคัน เนื่องจากการแพ้อาหาร แพ้ยาบางชนิด แพ้เครื่องสำอาง แพ้แมลง ผื่นคันตามก้น

วิธีใช้ ใช้ทาตามผิวหนังที่เป็นผื่นคัน และถูเล็กน้อยให้ยาซึมซาบลงไป ทาวันละ 1-2 ครั้ง หรือตามแพทย์สั่ง ในโรคผิวหนัง นอกจากให้ยาทาเฉพาะที่แล้ว ควรวิเคราะห์หาสาเหตุด้วย เมื่อพบเหตุแล้วก็รักษาเสียพร้อมกัน จะทำให้โรคหายเร็วขึ้น

คำเตือน ห้ามใช้ในผู้ที่แพ้ยานี้



11. Prednisolone (5 mg/tab)

MIMS 99th Edition 2005 page 195

I: RA, rheumatic fever, status asthmaticus, allergic & inflammatory conditions.

D: 1- 6 tab daily (individualized dosage).

CI: Acute superficial herpes simplex keratitis, systemic fungal infection, varicella.

SP: Hyperthyroidism, cirrhosis, ulcerative colitis, hypertension, CHF, peptic ulcer, DM. Childn & elderly.

AR: Insomnia, nervousness, increased appetite, indigestion, hirsutism, DM, cataract, glaucoma.

DI: Phenytoin, rifampin, barbiturates, salicylates.
US FDA Preg Cat, ♀ : C; D in 1st trimester (systemic route).



Antifungal agents

1. Daktarin oral gel 2 % (10 gm/pack)

Sig. apply tid pc.

C: Miconazole nitrate.

I: Curative & prophylactic treatment of candidosis of the oropharyngeal cavity & the GI tract.

D: Adult & childn ½ tsp qid. Infant ½ tsp bid.

CI: Liver dysfunction.

AR: Nausea, diarrhoea(long term).

DI: Increase the anticoagulant effect of coumarin derivatives, potentiates the effect of oral hypoglycemics, slows the metabolism of phenytoin & cyclosporin.

US FDA Preg Cat, ♀ : C

Note : store below 25 °C

MIMS 99th Edition 2005 page 359



2. Mycostatin™ oral suspension (100,000 units/ml) (12 ml/bottle)

Sig. dropped 1 ml into the mouth & held for some time before swallowing.

ชนิดน้ำแขวนตะกอน เป็นยาน้ำรสหอมหวาน มีไนสตาทิน 1 แสนหน่วย ต่อมล. เป็นยาปฏิชีวนะต่อต้านเชื้อราและยีสต์ โดยเฉพาะอย่างยิ่งเชื้อรา จำพวกแคนดิดา หรือ โมนิเลีย

ข้อบ่งใช้ ใช้รักษาโรคที่เกิดจากเชื้อราแคนดิดาของช่องปาก ทางเดินอาหาร ถ้าได้ สามารถใช้รักษาพร้อมกับยาปฏิชีวนะอื่นได้ ใช้ป้องกันการติดเชือรานในเด็กเกิดใหม่ กรณีที่ตรวจพบเชื้อราบริเวณช่องคลอดของมารดา

คำเตือน

1. ห้ามใช้ในผู้ที่แพ้ยานี้
2. ยานี้ อาจทำให้เกิดอันตรายต่อดับ ไต ได้
3. ยานี้มี alcohol ผสมอยู่ 1 % v/v ควรใช้ด้วยความระมัดระวัง



3. Nystin vaginal tab

Sig. 1 tab use as lozenge od-bid.

C : Nystatin 100,000 iu.

Diiodohydroxyquinoline 100 mg.

Benzalkonium Chloride 7 mg.

I : Vaginitis due to fungi, trichomonas, bacterial infections or nonspecific vaginitis.

D : 1 tab once-bid for 1 wk.

CI : Iodine intolerance.

AR : Local irritation. Iodism.

US FDA Preg Cat, (♀)

MIMS 99th Edition 2005 page 294



4. Ketoconazole (Ninazol®)

Sig. 1 tab orally od.

Each tablet contains : Ketoconazole 200 mg.

I: Systemic fungal infections, fungal infections of skin, hair & nails, candidiasis, pityriasis versicolor.

D : Adult 1 tablet (200 mg) once daily.

CI: Hepatic impairment.

SP: Pregnancy & lactation.

AR: Nausea; rashes, pruritus. Rarely, hepatotoxicity, gynaecomastia.

DI: Reduced absorption with anticholinergics, antacids & cimetidine.

US FDA Preg Cat, ♀ : C



Antiviral agents

1. Acyclovir cream (Vilerm®) 5 gm/pack Sig. apply to the lesion 5 times daily.

Each 100 g. contains Acyclovir 5 g.

Indications: Treatment of herpes simplex virus infections of the skin initial and recurrent genital herpes & herpes labialis.

Contraindications: Hypersensitivity or chemical intolerance to the components of the formulation.

Precautions & warnings

- For cutaneous use only. Do not use in the eye.



2. Acyclovir (Zovirax®) (200, 400, 800 mg/tab) ไม่มียาภายในคณะ - MIMS 99th Edition 2005 page 275 Sig Adult & child > 2 yr. Herpes simplex 1 tabs orally 5 times daily for 5 days. Herpes zoster 4 tabs 5 times daily for 7 days.

C: Acyclovir

D: Oral Adult & child > 2 yr. **Herpes simplex** 200 mg 5 times daily approx 4 hrly omitting the night dose for 5 days.

Herpes zoster (shingles) 800 mg 5 times daily approx 4 hrly omitting the night dose for 7 days.

Varicella (chickenpox) Adult 800 mg 5 times daily approx 4 hrly omitting the night dose for 7 days. Childn > 6 yr 800 mg qid for 5 days, 2-6 yr 400 mg qid for 5 days, < 2 yr 200 mg qid for 5 days. Childn Approx dosing: 20 mg/kg body wt qid.

SP: In pt receiving Zovirax IV at higher doses, specific care regarding renal function should be taken, particularly when pt are dehydrated or have any renal impairment.

AR: Skin rashes; GI effects. Rare: Neurological reactions(IV infusion).

DI: Increased mean ½-life & plasma conc with probenecid.

US FDA Preg Cat, ♀ : B

Mouthwashes

1. Sodium bicarbonate mouthwash (300 ml/bottle)

Sig. use as mouthwash tid pc.

Note : หมดอายุหลังเติมน้ำ 6 เดือน



2. Chlorhexidine 0.2 % mouthwash (240 ml/bottle)

Sig. use as mouthwash bid pc.

คำแนะนำ : ห้ามใช้ยานี้ติดต่อกัน นานเกิน 7 วัน



3. Hydrogenperoxide 1% solution mouthwash (240 ml/bottle)

Sig. use as mouthwash bid pc.



4. Difflam solution 0.15% (100mL/bottle)

Sig. mouth rinse 5-15 mL. for 30 seconds tid pc.

MIMS 99th Edition 2005 page 360

C: Benzydamine HCl

I: Sore throat, tonsillitis, radiation mucositis, aphthous ulcers, post-rosurgical & periodontal procedures.

D: Soln Gargle or rinse for 30 seconds, usually undiluted.
Adult 15 mL every 1½– 3 hr.
Childn 5-15 mL. Uninterrupted treatment should not be > 7 days.

SP: Do not swallow. Hepatic or renal impairment; pregnancy, Childn < 6 yr.

AR: Oral numbness; dryness or thirst, tingling, warm feeling in mouth, altered sense of taste.

US FDA Preg Cat, (♀)



5. Sodium fluoride mouthwash 0.055% (120 ml/bottle)



Anticonvulsants

1. Carbamazepine (Tegretol®) (200 mg/tab) Sig. 100 - 1,600 mg/day orally.

MIMS 99th Edition 2005 page 165

I: Epilepsy (except petit mal). Alcohol withdrawal syndrome. Trigeminal neuralgia, idiopathic glossopharyngeal neuralgia. Diabetic neuropathy. Diabetes insipidus. Mania, prophylaxis in manic depressive disorders.

D: Adult 100 - 1600 mg/day depending on indication & severity of illness.
Children 10-20 mg/kg/day in divided doses.

CI: AV block. Concomitant use of MOIs. History of bone marrow depression or acute intermittent porphyria.

SP: Pregnancy, lactation. Initial & periodic complete blood counts, liver function tests & urine analysis. Monitoring of plasma levels. History of cardiac, hepatic or renal damage & adverse haematological reactions. Road / machinery users. Intraocular pressure, latent psychosis, confusion, agitation. Mixed seizure disorders. Elderly patients. Abrupt discontinuation of treatment.

AR: Frequent: dizziness, ataxia, mild allergic skin reactions, mild leucopenia. Occasionally: diplopia, liver enzyme elevations, vomiting, thrombocytopenia, hyponatremia, accommodation disorders. Rare: exfoliative dermatitis, Stevens-Johnson syndrome, jaundice, hepatitis, delayed multi-organ hypersensitivity disorders, SLE-like syndrome. Isolated cases: neuritis, hallucinations, agitation, depression, Lyell' syndrome, blood dyscrasias, anaphylactic reaction, AV-block, congestive heart failure, thromboembolism, osteomalacia, renal dysfunction, interstitial nephritis, sexual disturbances.

DI: Carbamazepine is a powerful enzyme inducer. Clinically important interaction with other drugs/substances(including OCs alcohol) are common.

P/P: Tab 200,400 mg Syr 100mg/5ml × 250 ml

US FDA Preg Cat, ♀ : D



2. **Phenytoin (Ditoin) (100 mg/cap)**

Sig. 100 mg orally tid.

C: Phenytoin Na.

I: Psychomotor epilepsy, migraine & trigeminal neuralgia.

D: Adult 100 mg tid increased to 200 mg tid if necessary.
Children 30 mg tid.

CI: Pregnancy.

SP: Hepatic disease, avoid abrupt withdrawal; lactation.

AR: Nystagmus, ataxia, slurred speech, confusion, dizziness, gingival hyperplasia, hirsutism, mobiliform rash, rickets, osteomalacia, lupus erythematosus syndrome, leukopenia, thrombocytopenia, pancytopenia, granulocytopenia.

DI: Metabolism enhanced by barbiturates & inhibited by chloramphenicol, dicoumarol anticoagulants, disulfiram, phenylbutazone, INH, sulfafenazole, sulthiame, valproic acid, cimetidine & sulfonamides.

US FDA Preg Cat, ♀ : D

MIMS 99th Edition 2005 page 164



3. **Clonazepam (Rivotril®) (0.5 mg/tab)**

Sig. 1 tab orally od or bid or tid.

C: Clonazepam.

I: Typical or atypical petit mal epilepsies, grand mal, status epilepticus in all its clinical forms.

D: Adult initially 1-1.5 mg/day tid. Maintenance: 3-6 mg/day. Childn 10-16 yr initially 1-1.5 mg/day bid-tid. Maintenance: 3-6 mg/day.

SP: Pt with spinal or cerebella ataxia, in the event of acute alcohol or drug intoxication; severe liver damage. COPD. May modify Pt' reaction. Abrupt withdrawal. History of depression & suicide attempts. Simutaneous use of other antiepileptics. Pregnancy & lactation.

AR: Tiredness, sleepiness, lassitude, muscular hypotonia, muscle weakness, dizziness, light headedness, ataxia, slowed reaction, poor conc., restlessness, confusion, disorientation; anterograde amnesia, depression. Increased production of saliva or of bronchial secretions in infants & young childn.

US FDA Preg Cat, ♀ : D

MIMS 99th Edition 2005 page 165



4. **Gabapentin (Neurontin®) (100, 300, 400 mg/cap)** ไม่มียาภายในคณะ - MIMS 99th Edition 2005 page 164

Sig. adults initially 900 mg/day orally given as 3 divided dose. May be initiated with 300 mg tid on day 1 or with 3-day titration by giving 300 mg on day 1, 300 mg bid on day 2, 300 mg tid on day 3. Titrated if necessary up to max of 3,600 mg/day.

I: Neuropathic pain in adults \geq 18 yr.

Muscle Relaxants

1. **Baclofen (Baclosal®) (10 mg/tab)**

MIMS 99th Edition 2005 page 181

Sig. 15-80 mg/day orally.

I: Muscle relaxation for muscle spasticity of spinal & cerebral origin including spasticity of the skeletal muscles in multiple sclerosis.

D: Initially 5 mg tid. Maintenance: 30-80 mg/day.

SP: Psychiatric disorders, epilepsy, pregnancy, CV or cerebrovascular insufficiency. May impair ability to drive or operate machinery.

AR: Sedation, drowsiness, nausea.

DI: Antidepressants, levodopa, antihypertensives.

US FDA Preg Cat, ♀ : C



Antidepressants

1. **Amitriptyline HCl. (Tripta®) (10 mg/tab)**

MIMS 99th Edition 2005 page 171

Sig 1 tab orally od.

I: Mental depression.

D : Initial: 75-150 mg daily in divided doses.

Maintenance: 50-100 mg daily in divided doses.

CI: Immediate recovery phase after MI, heart block. Mania. MAOI therapy.

SP: CV diseases. Hyperthyroidism. Impaired liver function. History of epilepsy, suicidal tendencies. Narrow-angle glaucoma, urinary retention, prostatic hypertrophy. Diabetes. Elderly, childn < 5 yr.

AR: Dry mouth, constipation, urinary retention, blurred vision, narrow-angle glaucoma, impotence, drowsiness, dizziness, sweating, weakness, fatigue, epileptiform seizures, orthostatic hypotension, palpitation, tachycardia, cardiac arrhythmia, extrapyramidal symptoms, endocrine effects.

DI: Barbiturates, phenytoin, carbamazepine, phenothiazines, cimetidine, estrogen, OC, alcohol, sympathomimetic amines, debrisoquine, guanethidine, clonidine, bethanidine.

US FDA Preg Cat, ♀ : C



Others agents

1. Glycerin borax (10 ml/bottle) Sig. apply 3-5 times daily.

Note : shake well before use.



2. Solcoseryl dental adhesive paste (5 gm/pack) Sig. apply to dried lesions 3-5 times daily.

MIMS 99th Edition 2005 page 361

C: Protein-free haemodialysate 5%, polidocanol 1%

I: Lesions of the oral mucosa, gums & lips eg aphthae, rhagades, herpes labialis, gingivitis, parodontitis, stomatitis; denture pressure sores, teething pains; dressing after scaling, curettage & dental extractions; collocation of immediate dentures.

US FDA Preg Cat, (♀)



3 . Sore mouth gel (5 gm/pack) Sig. apply 3 hrly as required.

MIMS 99th Edition 2005 page 361

C: Per 100 g lignocaine base 0.6 g, cetylpyridinium Cl 0.02 g, menthol 0.06 g, eucalyptol 0.1 mL.

I: Relief from pain caused by ulcers, lesions & abrasions of mouth & lips.

D: Apply 3 hrly as required.

US FDA Preg Cat, (♀)



4.. Kamistad®-Gel (10 gm/pack)
Sig. apply tid pc.

MIMS 99th Edition 2005 page 360

C : 1 gm of gel contains : lidocaine hydrochloride 1 H₂O, 20 mg; thymol; chamomile flower tincture(1:5.5), 200 mg; and Benzalkonium chloride, 1 mg.

I : For painful, inflamed conditions of the mucous membrane of the mouth and lips. For denture wearers, for rubbing into the gums at pressure points.

US FDA Preg Cat, (♀)



5. Kamillosan-M (Spray 15 mL)

ไม่มียาภายในคณะ - MIMS 99th Edition 2005 page 360

C: Per 100 mL Standardised chamomile extr 370.5 mg, essential oil 100 mg%, chamazulene 3 mg%.

I: Inflammatory affections of buccal & pharyngeal cavity, bad breath, parodontosis, acute gingivitis, pain after tooth extraction, mucosal irritation, tonsillary angina, canker sores.

D: Spray tid.

SP: Do not spray directly into the nose.

US FDA Preg Cat, (♀)

♀ represents U.S.FDA Pregnancy Category. Refer to Pg A10 for definition of Categories A, B, C, D & X. (♀) represents products which currently do not have a pregnancy category or where the pregnancy category is not applicable. No representations on the safety of a drug in pregnancy can be made for products without a pregnancy category.

Vitamins & Minerals

1. Vitamin B1-B6-B12

Sig. 1 tab orally od.

Each tablet contain

Vitamin B1 Mononitrate	100 mg
Vitamin B6	7.5 mg
Vitamin B12	75.0 mcg

Use : Prevention & treatment vitamin B1-B6-B12 Deficiency.

Dose : 1 tab orally daily or as directed by the physician



2. Vitamin B complex

Sig. 1-2 tab orally tid.

Each tablet contain

Vitamin B1 Mononitrate	5 mg
Vitamin B2	2 mg
Vitamin B6	2 mg
Nicotinamide	20 mg

Use : Prevention & treatment vitamin B deficiency

Dose : 1-2 tab orally tid.



3. Vitamin C


Sig. 1-2 tab orally od-bid.

Each tablet contain

Vitamin C (Ascorbic acid)	100 mg.
---------------------------	---------

Indications : Use : Vitamin C deficiency, antiscorbutic.

Dose : 1-2 tab orally od or bid.



4. Folic acid 5 mg/tab (Folivit)

MIMS 99th Edition 2005 page 336

Sig. 1-4 tab orally daily.

I: Megaloblastic anemia due to a deficiency of folic acid (as may be seen in tropical or non-tropical sprue) & in anemia of nutritional origin, pregnancy, infancy or childhood.

D: 1-4 tab daily.



5. FBC

MIMS 99th Edition 2005 page 335

Sig. 1 tab orally bid-tid.

C: Fe fumarate	200 mg
Vit B1	2 mg
Vit B2	2 mg
Vit B12	5 mcg
Vit C	20 mg
Niacin	10 mg
Folic acid	100 mcg
Ca phosphate tribasic	100 mg.

I: Anemia

D: 1 tab bid-tid



6. Ferli-6

MIMS 99th Edition 2005 page 335

Sig. 1 tab orally bid-tid.

C: Fe fumarate 200 mg, folic acid 0.5 mg, pyridoxine HCl 5 mg.

I: Anemia, Vit & mineral supplement.

D: 1 tab bid-tid



Vitamins & Minerals

ไม่มียาในคณะ

Centrum (Zuellig)

MIMS 99th Edition 2005 page 330

C :	Vit A (as acetate)	2,000	iu
	Vit B ₆	3	mg
	Vit B ₁₂	9	mcg
	Vit C	90	mg
	Vit D	400	iu
	Vit E	30	iu
	B-carotene(equiv to Vit A)	3,000	iu
	Biotin	45	mcg
	Folic acid	400	mcg
	Niacinamide	40	mg
	Pantothenic acid	10	mg
	Riboflavin	3.2	mg
	Thiamine	2.25	mg
	Ca	175	mg
	Cl	36	mg
	Chromium	25	mcg
	Copper	2	mg
	Iodine	0.15	mg
	Fe	10	mg
	Mg	100	mg
	Manganese	5	mg
	Molybdenum	25	mcg
	Nickel	5	mcg
	Phosphorus	125	mg
	K	40	mg
	Selenium	25	mcg
	Silica	10	mcg
	Tin	10	mcg
	Vanadium	10	mcg
	Zn	15	mg

I: Vit & mineral deficiency due to illness, poor appetite &/or drug-induced malabsorption.

D: Adult 1 tab daily.

Neurobion

MIMS 99th Edition 2005 page 325

C : Per tab			
	Vit B ₁	100	mg
	Vit B ₆	200	mg
	Vit B ₁₂	200	mcg
Per amp			
	Vit B ₁	100	mg
	Vit B ₆	100	mg
	vit B ₁₂	1	mg

I: Toxic & nontoxic neuritis; diabetic & alcoholic neuropathy, drug-induced neuropathy & depression; toxæmia of pregnancy; anorexia, neurasthenia; general tonic during pregnancy, in convalescence & growth. **Tab** For moderate cases or as follow-up therapy for a course of inj. **Inj** For severe cases.

D: Tab 1 tab tid. Inj 1 amp/day IM. After resolution of acute symptoms: 1 amp IM 2-3 times wkly.

SP: Prolonged treatment.

DI: High vit B6 does decrease action of L-dopa.

Stresstabs 600

MIMS 99th Edition 2005 page 325

C :	Vit B ₁	15	mg
	Vit B ₂	15	mg
	Vit B ₆	5	mg
	Vit B ₁₂	12	mcg
	Vit C	600	mg
	Vit E	30	iu
	Folic acid	400	mcg
	Niacinamide	100	mg
	Biotin	45	mcg
	Pantothenic acid	20	mg

I: Treatment of water-soluble vit deficiency, during serious illness, burns & surgery, fracture & trauma, alcoholism & drug abuse, wound healing.

D: Adult 1 tab daily.

Stresstabs 600 + Iron

MIMS 99th Edition 2005 page 332

C :	Vit B ₁	15	mg
	Vit B ₂	10	mg
	Vit B ₆	5	mg
	Vit B ₁₂	12	mcg
	Vit C	600	mg
	Vit E	30	iu
	Nicotinamide	100	mg
	Pantothenic acid	20	mg
	Biotin	45	mcg
	Folic acid	400	mcg
	Fe	27	mg

I: Water soluble vit deficiency. Serious illness, burns & surgery, fracture & trauma, alcoholism & drug abuse, wound healing.

D: Adult 1 tab daily.

Stresstabs 600 + Zinc

MIMS 99th Edition 2005 page 332

C :	Cupric oxide	3	mg
	Vit B ₁	15	mg
	Vit B ₂	10	mg
	Vit B ₆	5	mg
	Vit B ₁₂	12	mcg
	Vit C	600	mg
	Vit E	30	iu
	Nicotinamide	100	mg
	Pantothenic acid	20	mg
	Biotin	45	mcg
	Folic acid	400	mcg
	Zn	23.9	mg

I : Water soluble vit deficiency. Serious illness, burns & surgery, fracture & trauma, alcoholism & drug abuse, wound healing.

D : Adult 1 tab daily.

Tribesian

MIMS 99th Edition 2005 page 325

C :	Vit B ₁ mononitrate	100	mg
	Vit B ₆ HCl	5	mg
	Vit B ₁₂	50	mcg

I : Neuralgias, migraine, neuritis. Anaemic conditions.

D : 1 tab daily.

Z-Bec

MIMS 99th Edition 2005 page 333

C :	Vit B ₁	15	mg
	Vit B ₂	10.2	mg
	Vit B ₆	10	mg
	Vit B ₁₂	6	mcg
	Vit C	600	mg
	Vit E	45	iu
	Niacin	100	mg
	Pantothenic acid	25	mg
	Zn	22.5	mg

I : Dietary supplement.

D: Adult & childn > 12 yr 1 tab daily.

SP: Not intended for the treatment of pernicious anemia.